



It is wrong to treat pain in CSF hypotension headache (post-lumbar puncture headache) to allow the patient to walk!

Marcelo Moraes Valença¹ , Matheus Salerno Sousa² , Marcelo Andrade Valença¹ ,
Déborah Emmily de Carvalho³ , Alberto Henrique Torres Trindade da Silva³ , Juliana Ramos de Andrade¹ 

¹Federal University of Pernambuco, Recife, Pernambuco, Brazil

²Franca Municipal University Center (Uni-Facef), Franca, São Paulo, Brazil

³University of Pernambuco, Recife, Pernambuco, Brazil



Marcelo Moraes Valença
mmvalenca@yahoo.com.br

Edited by:

Mario Fernando Prieto Peres

A secondary headache, common in daily practice, is the post-dural puncture headache, which is very frequent in spinal anesthesia. Some patients in the immediate postoperative period, when he or she got out of bed, report the appearance of pain of moderate to severe intensity that improves when he or she returns to the lying position.¹⁻³ Valença and coworkers², in an experiment with fragments of human dura mater obtained from cadaver donors of both sexes, evaluated why women are more susceptible than men to development post-dural puncture headache (PDPH). The experiment demonstrated that the dura mater has mechanisms that provide spontaneous occlusion of the orifice caused by the entrance of the needle, even when placed isolated and fixed an acrylic apparatus simulating the human spine in the vertical position, under the pressure of 40 cm H₂O. This particularity observed in cadaver dura mater varies between individuals and also between different parts of the dura mater of the same individual. This may explain why some individuals have post-dural puncture headache and others do not. The experimental data revealed that dura mater fragments obtained from women leaked more fluid, and the orifice remained opened longer when punctured when compared with men.

The triggering of pain during biological events that have the potential to be life-threatening is a crucial mechanism that is often overlooked. Numerous published articles are bringing different alternatives in an attempt to treat a postural headache that is pathognomonic of cerebrospinal fluid (CSF) hypotension (Table 1).⁴⁻²³ Even so, one of the most used drugs to prevent and treat post-lumbar puncture headache, caffeine, had its analgesic action questioned in a publication of notable reputation due to being insufficiently supported by the available pharmacological and clinical proofs.²⁴

From a practical point of view, it is worth remembering that normal CSF pressure is not uncommon in patients with spontaneous intracranial hypotension and, thus, the lack of a low opening pressure does not exclude this condition.²⁵ And spontaneous intracranial hypotension may be encountered in patients with new-onset daily persistent headache.²⁶

There are several causes of CSF hypotension,²⁷⁻³⁶ the most frequent after the spinal anesthesia procedure.³⁷

Keywords:

Post-Dural Puncture Headache
Intracranial Hypotension
Blood Patch
Epidural
Headache
Treatment



In the intracranial space, there are three compartments: CSF, brain parenchyma, and blood. With the exception of the brain parenchyma, which is practically inelastic, there is a considerable variation during the cardiac cycle in the volumes of CSF and blood. These intracranial volumes maintain an inversely directional relationship. When there is an increase in the intracranial volume of blood, the CSF must leave the cavity towards the spinal dural sac to allow blood to enter the intracranial cavity. Cerebrospinal fluid leaves and enters through the foramen magnum about 70 times per minute in a resting individual.

Table 1. Treatment alternatives in post-dural puncture headache.

Modified sphenopalatine ganglion block ^{4,8}
Sphenopalatine block with lidocaine spray ⁶
Intranasal lidocaine atomization ⁵
Epidural blood patch ³⁸⁻⁵³
Synacthen depot ¹⁵
Caffeine ^{14,23,24}
Sumatriptan ^{9,20,22}
Naratriptan ⁹
Aminophylline/Theophylline ¹⁰
Piritramid ¹²
Ergotamine ¹⁴
Gabapentin ¹⁴
Tramadol ¹¹
Paracetamol ¹²
Cosyntropin ^{7,18,21}
Tetracosactin ¹⁶

In a reduced volume of the CSF, in the case of fistula with loss of CSF, the pain is caused by the exit of the CSF from the intracranial space into the spinal dural sac. In order to keep the intracranial volume constant with the three mentioned compartments (i.e., brain parenchyma, CSF, and blood), there is vasodilation, mainly of the venous sinuses and large veins, as these vessels are quite innervated, the individual feels an intense headache when raising the head in relation to the trunk. By the force of gravity, this position facilitates the exit of the already reduced volume of CSF from the intracranial cavity to the dural sac. Another explanation for postural headache is the descent of the cerebellar tonsils through the foramen magnum, as in the case of a secondary Chiari,⁵⁴ if that name is correct for this diagnosis.

Cerebrospinal fluid hypotension due to a reduced volume of CSF, as in the example of a fistula, can precipitate complications such as bleeding in the subdural space. It is not uncommon for chronic subdural hematomas to appear after

the patient has undergone spinal anesthesia or after a lumbar puncture with CSF removal.⁵⁵ Cerebrospinal fluid hypotension may be a cause of cortical vein thrombosis.⁵⁶ In addition, cerebellar hemorrhage may also be found as a manifestation of spontaneous intracranial hypotension.²⁶

The authors, therefore, is not against the treatment of post-dural puncture headache or the headache associated with cerebrospinal fluid hypotension from other causes as long as it treats the cause of cerebrospinal fluid fistula, as in the example of epidural blood patch.^{39-44, 57} Also, remember that epidural blood patch is not an uncomplicated procedure (e.g., arachnoiditis, seizures and respiratory distress, pneumocephalus, intracranial subarachnoid hemorrhage, hydrocephalus, local infection and sepsis complication, and spinal subdural hematoma)^{38, 45-51} and should be planned and indicated when conservative measures, including rest in the horizontal position and hydration, have failed.^{12, 52, 57} In most cases, the headache disappears within a week.¹ In the case of prolonged horizontal decubitus, measures to prevent deep venous thrombosis should be taken. Other forms of CSF fistula treatment have also been published.^{53, 58-61}

Thus, we conclude that it is wrong to treat pain in patients with the postural headache associated with a CSF hypotension with the intention to allow the patient to walk because it favors the occurrence of intracranial complications that can lead to the patient's death. The pain makes the patient remain in the horizontal decubitus and, in this way, avoids the exit of the CSF to the spinal dural sac, causing a substantial intracranial hemodynamic disturbance with hemorrhagic or thrombotic repercussions.

Conflict of interest: no conflict of interest to declare

Authors' contribution: MMV – writing: original draft, review & editing, visualization, supervision; JRA – writing: review & editing, visualization, supervision; MSS - writing: editing, table and references; MAV, DEC, AHTTS - writing: editing, table and references

Marcelo Moraes Valença
<https://orcid.org/0000-0003-0678-3782>
 Matheus Salerno Sousa
<https://orcid.org/0000-0002-2335-8019>
 Marcelo Andrade Valença
<https://orcid.org/0000-0002-0824-0928>
 Déborah Emmily de Carvalho
<https://orcid.org/0000-0002-8068-3598>



Alberto Henrique Torres Trindade da Silva
<https://orcid.org/0000-0003-2356-4426>
 Juliana Ramos de Andrade
<https://orcid.org/0000-0002-5445-8872>

References

- Amorim JA, Gomes de Barros MV and Valenca MM. **Post-dural(post-lumbar)punctureheadache:riskfactors and clinical features.** *Cephalalgia* 2012;32(12):916-923 Doi:10.1177/0333102412453951
- Valenca MM, Amorim JA and Moura TP. **Why don't all individuals who undergo dura mater/arachnoid puncture develop postdural puncture headache?** *Anesth Pain Med* 2012;1(3):207-209 Doi:10.5812/kowsar.22287523.3616
- Amorim JA and Valenca MM. **Postdural puncture headache is a risk factor for new postdural puncture headache.** *Cephalalgia* 2008;28(1):5-8 Doi:10.1111/j.1468-2982.2007.01454.x
- Bhargava T, Kumar A, Rastogi A, Srivastava D and Singh TK. **A Simple Modification of Sphenopalatine Ganglion Block for Treatment of Postdural Puncture Headache: A Case Series.** *Anesth Essays Res* 2021;15(1):143-145 Doi:10.4103/aer.aer_67_21
- Siegler BH, Gruss M, Oehler B, Kessler J, Fluhr H, Weis C, . . . Weigand MA. **[Intranasal lidocaine atomization as novel and noninvasive treatment option for postdural puncture headache : Two case reports from obstetric anesthesiology].** *Anaesthesist* 2021;70(5):392-397 Doi:10.1007/s00101-020-00900-9
- Lopez T, Sastre JA and Gomez-Rios MA. **Sphenopalatine block with lidocaine spray for treatment of obstetric postdural puncture headache.** *J Clin Anesth* 2021;68(110069 Doi:10.1016/j.jclinane.2020.110069
- Zapapas MK, Gralla J, Tong S and Eisdorfer S. **Cosyntropin for the Treatment of Refractory Postdural Puncture Headache in Pediatric Patients: A Retrospective Review.** *Clin J Pain* 2020;36(3):213-218 Doi:10.1097/AJP.0000000000000784
- Liu CW and Lim H. **Sphenopalatine ganglion block for the treatment of postdural puncture headache.** *Comment on Br J Anaesth* 2020 124: 739-47. *Br J Anaesth* 2020;125(4):e359 Doi:10.1016/j.bja.2020.05.043
- Botros JM and Sayed AM. **Comparison between the Effects of Sumatriptan Versus Naratriptan in the Treatment of Postdural Puncture Headache in Obstetric Patients: A Randomized Controlled Trial.** *Anesth Essays Res* 2019;13(2):376-382 Doi:10.4103/aer.AER_17_19
- Wu C, Guan D, Ren M, Ma Z, Wan C, Cui Y, . . . Xie N. **Aminophylline for treatment of postdural puncture headache: A randomized clinical trial.** *Neurology* 2018;90(17):e1523-e1529 Doi:10.1212/WNL.0000000000005351
- Stephenson LL, Varness DC, Schroeder KM and Ford MP. **Tramadol for postdural puncture headache treatment in a pediatric patient following failed blood patch.** *J Clin Anesth* 2012;24(2):171-172 Doi:10.1016/j.jclinane.2011.03.009
- Marcus HE, Fabian A, Dagtekin O, Schier R, Krep H, Bottiger BW, . . . Petzke F. **Pain, postdural puncture headache, nausea, and pruritus after cesarean delivery: a survey of prophylaxis and treatment.** *Minerva Anesthesiol* 2011;77(11):1043-1049 Doi:R02116532 [pii]
- Kleine-Bruggeney M, Kranke P and Stamer UM. **[Prophylaxis and therapy of postdural puncture headache--a critical evaluation of treatment options].** *Anesthesiol Intensivmed Notfallmed Schmerzther* 2011;46(7-8):516-524 Doi:10.1055/s-0031-1284471
- Erol DD. **The analgesic and antiemetic efficacy of gabapentin or ergotamine/caffeine for the treatment of postdural puncture headache.** *Adv Med Sci* 2011;56(1):25-29 Doi:10.2478/v10039-011-0009-z
- Rucklidge MW, Yentis SM and Paech MJ. **Synacthen Depot for the treatment of postdural puncture headache.** *Anaesthesia* 2004;59(2):138-141 Doi:10.1111/j.1365-2044.2004.03573.x
- Canovas L, Barros C, Gomez A, Castro M and Castro A. **Use of intravenous tetracosactin in the treatment of postdural puncture headache: our experience in forty cases.** *Anesth Analg* 2002;94(5):1369 Doi:10.1097/00000539-200205000-00069
- Benzon HT and Wong CA. **Postdural puncture headache: mechanisms, treatment, and prevention.** *Reg Anesth Pain Med* 2001;26(4):293-295 Doi:10.1053/rapm.2001.23670
- Carter BL and Pasupuleti R. **Use of intravenous cosyntropin in the treatment of postdural puncture headache.** *Anesthesiology* 2000;92(1):272-274 Doi:10.1097/00000542-200001000-00043
- Van de Velde M, Corneillie M, Vanacker B, Stevens E, Verhaeghe J, Van Assche A and Vandermeersch E. **Treatment for postdural puncture headache associated with late postpartum eclampsia.** *Acta Anaesthesiol Belg* 1999;50(2):99-102, <http://www.ncbi.nlm.nih.gov/pubmed/10418650>



20. Sprigge JS. **The use of sumatriptan in the treatment of postdural puncture headache after accidental lumbar puncture complicated a blood patch procedure.** *Anaesthesia* 1999;54(1):95-96 Doi:10.1046/j.1365-2044.1999.0759q.x
21. Kshatri AM and Foster PA. **Adrenocorticotrophic hormone infusion as a novel treatment for postdural puncture headache.** *Reg Anesth* 1997;22(5):432-434 Doi:10.1016/s1098-7339(97)80029-6
22. Hodgson C and Roitberg-Henry A. **The use of sumatriptan in the treatment of postdural puncture headache.** *Anaesthesia* 1997;52(8):808, <https://www.ncbi.nlm.nih.gov/pubmed/9291788>
23. Baumgarten RK. **Should caffeine become the first-line treatment for postdural puncture headache?** *Anesth Analg* 1987;66(9):913-914, <https://www.ncbi.nlm.nih.gov/pubmed/3619102>
24. Halker RB, Demaerschalk BM, Wellik KE, Wingerchuk DM, Rubin DI, Crum BA and Dodick DW. **Caffeine for the prevention and treatment of postdural puncture headache: debunking the myth.** *Neurologist* 2007;13(5):323-327 Doi:10.1097/NRL.0b013e318145480f
25. Kranz PG, Tanpitukpongse TP, Choudhury KR, Amrhein TJ and Gray L. **How common is normal cerebrospinal fluid pressure in spontaneous intracranial hypotension?** *Cephalgia* 2016;36(13):1209-1217 Doi:10.1177/0333102415623071
26. Schievink WI, Maya MM and Nuno M. **Chronic cerebellar hemorrhage in spontaneous intracranial hypotension: association with ventral spinal cerebrospinal fluid leaks: clinical article.** *J Neurosurg Spine* 2011;15(4):433-440 Doi:10.3171/2011.5.SPINE10890
27. Schievink WI, Michael LM, 2nd, Maya M, Klimo P, Jr. and Elijovich L. **Spontaneous Intracranial Hypotension Due to Skull-Base Cerebrospinal Fluid Leak.** *Ann Neurol* 2021;90(3):514-516 Doi:10.1002/ana.26175
28. Villamil F, Ruella M, Perez A, Millar Vernetti P, Paday Formenti ME, Acosta JN and Goicochea MT. **Traumatic vs Spontaneous Cerebrospinal Fluid Hypotension Headache: Our experience in a series of 137 cases.** *Clin Neurol Neurosurg* 2020;198:106140 Doi:10.1016/j.clineuro.2020.106140
29. Shah VN and Dillon WP. **Spontaneous Intracranial Hypotension With Brain Sagging Attributable to a Cerebrospinal Fluid-Venous Fistula.** *JAMA Neurol* 2020;77(10):1320-1321 Doi:10.1001/jamaneurol.2020.2372
30. Sanchez Ortega JF, Pinilla Arias D, Vazquez Miguez A and Calatayud Perez JB. **Intracranial hypotension secondary to spontaneous spinal cerebrospinal fluid fistula: Three case reports.** *Neurocirugia (Astur : Engl Ed)* 2020;Doi:10.1016/j.neucir.2020.07.006
31. Takai K, Niimura M, Hongo H, Umekawa M, Teranishi A, Kayahara T and Taniguchi M. **Disturbed Consciousness and Coma: Diagnosis and Management of Intracranial Hypotension Caused by a Spinal Cerebrospinal Fluid Leak.** *World Neurosurg* 2019;121(e700-e711 Doi:10.1016/j.wneu.2018.09.193
32. Dobrocky T, Grunder L, Breiding PS, Branca M, Limacher A, Mosimann PJ, . . . Piechowiak El. **Assessing Spinal Cerebrospinal Fluid Leaks in Spontaneous Intracranial Hypotension With a Scoring System Based on Brain Magnetic Resonance Imaging Findings.** *JAMA Neurol* 2019;76(5):580-587 Doi:10.1001/jamaneurol.2018.4921
33. Chan SM, Chodakiewitz YG, Maya MM, Schievink WI and Moser FG. **Intracranial Hypotension and Cerebrospinal Fluid Leak.** *Neuroimaging Clin N Am* 2019;29(2):213-226 Doi:10.1016/j.nic.2019.01.002
34. Silva AEJ, Pavan P, Oshima MM, Cardoso TM and Reis F. **Intracranial hypotension secondary to spontaneous spinal cerebrospinal fluid leaks.** *Arq Neuropsiquiatr* 2017;75(3):201-202 Doi:10.1590/0004-282X20160183
35. Tu A, Creedon K and Sahjipaul R. **Iatrogenic cerebrospinal fluid leak and intracranial hypotension after gynecological surgery.** *J Neurosurg Spine* 2014;21(3):450-453 Doi:10.3171/2014.5.SPINE13746
36. Liao YJ, Dillon WP, Chin CT, McDermott MW and Horton JC. **Intracranial hypotension caused by leakage of cerebrospinal fluid from the thecal sac after lumboperitoneal shunt placement. Case report.** *J Neurosurg* 2007;107(1):173-177 Doi:10.3171/JNS-07/07/0173
37. Ljubisavljevic S. **Postdural puncture headache as a complication of lumbar puncture: clinical manifestations, pathophysiology, and treatment.** *Neurol Sci* 2020;41(12):3563-3568 Doi:10.1007/s10072-020-04757-z
38. Ferrante E, Rubino F and Porrinis L. **Pneumocephalus: a rare complication of epidural catheter placement during epidural blood patch.** *Headache* 2014;54(3):539-540 Doi:10.1111/head.12235
39. Takai K and Taniguchi M. **Targeted Epidural Blood Patch Under O-Arm-Guided Stereotactic Navigation in Patients with Intracranial Hypotension Associated with a Spinal Cerebrospinal Fluid Leak and Ventral Dural Defect.** *World Neurosurg* 2017;107(351-357



- Doi:10.1016/j.wneu.2017.07.168
40. Faltings L, Kulason KO, Du V, Schneider JR, Chakraborty S, Kwan K, . . . Boockvar J. **Early Epidural Blood Patch to Treat Intracranial Hypotension after Iatrogenic Cerebrospinal Fluid Leakage from Lumbar Tubular Microdiscectomy.** *Cureus* 2018;10(11):e3633 Doi:10.7759/cureus.3633
 41. Wang E and Wang D. **Successful treatment of spontaneous intracranial hypotension due to prominent cervical cerebrospinal fluid leak with cervical epidural blood patch.** *Pain Med* 2015;16(5):1013-1018 Doi:10.1111/pme.12418
 42. Tanweer O, Kalhorn SP, Snell JT, Wilson TA, Lieber BA, Agarwal N, . . . Sutin KM. **Epidural Blood Patch Performed for Severe Intracranial Hypotension Following Lumbar Cerebrospinal Fluid Drainage for Intracranial Aneurysm Surgery. Retrospective Series and Literature Review.** *J Cerebrovasc Endovasc Neurosurg* 2015;17(4):318-323 Doi:10.7461/jcen.2015.17.4.318
 43. Chuang YS, Ju DT, Chiu TH, Huang YH, Cherng CH and Wu ZF. **Thoracic epidural blood patch with high volume blood for cerebrospinal fluid leakage of cervical spine (C2-3) complicated with spontaneous intracranial hypotension.** *Acta Anaesthesiol Taiwan* 2015;53(3):112-113 Doi:10.1016/j.aat.2015.06.003
 44. Mustafa R, Barman RA, Kurian EB, Eldrige J, Frank R, Kisson N, . . . Olatoye O. **Characterizing the need for multiple epidural blood patches in the treatment of postdural puncture headache: an observational study.** *Reg Anesth Pain Med* 2021;46(12):1110-1112 Doi:10.1136/rapm-2021-102692
 45. Seemiller J, Challagundla S, Taylor T and Zand R. **Intrathecal blood injection: a case report of a rare complication of an epidural blood patch.** *BMC Neurol* 2020;20(1):187 Doi:10.1186/s12883-020-01763-8
 46. Chazen JL and Amrhein TJ. **Arachnoiditis following epidural blood patch-An avoidable rare complication due to blind technique.** *Headache* 2021;61(6):972-973 Doi:10.1111/head.14145
 47. Hasiloglu ZI, Albayram S, Ozer H, Olgun DC, Selcuk H and Kaynar MY. **Cranial subarachnoid hemorrhage as an unusual complication of epidural blood patch.** *Clin Neurol Neurosurg* 2011;113(8):689-692 Doi:10.1016/j.clineuro.2011.04.005
 48. Al Maach N, Vogels OJ, Bollen TL and Wessels PH. **Arachnoiditis and communicating hydrocephalus as a complication of epidural blood patch.** *J Neurol* 2010;257(4):672-673 Doi:10.1007/s00415-009-5423-9
 49. Anwari JS. **Epidural blood patch (EBP) and septic complication.** *Can J Anaesth* 2000;47(3):289-290 Doi:10.1007/BF03018931
 50. Tekkok IH, Carter DA and Brinker R. **Spinal subdural haematoma as a complication of immediate epidural blood patch.** *Can J Anaesth* 1996;43(3):306-309 Doi:10.1007/BF03011749
 51. Reynolds AF, Jr., Hameroff SR, Blitt CD and Roberts WL. **Spinal subdural epiarachnoid hematoma: a complication of a novel epidural blood patch technique.** *Anesth Analg* 1980;59(9):702-703, <https://www.ncbi.nlm.nih.gov/pubmed/7191232>
 52. Katz D and Beilin Y. **Review of the Alternatives to Epidural Blood Patch for Treatment of Postdural Puncture Headache in the Parturient.** *Anesth Analg* 2017;124(4):1219-1228 Doi:10.1213/ANE.0000000000001840
 53. Wong K and Monroe BR. **Successful Treatment of Postdural Puncture Headache Using Epidural Fibrin Glue Patch after Persistent Failure of Epidural Blood Patches.** *Pain Pract* 2017;17(7):956-960 Doi:10.1111/papr.12541
 54. Yoshimoto S, Takai K, Takahashi K, Yasui T and Taniguchi M. **Intracranial hypotension and hypertension: reversible Chiari malformation due to dynamic cerebrospinal fluid abnormalities in Gorham-Stout disease. Case report.** *J Neurosurg Pediatr* 2018;22(5):508-512 Doi:10.3171/2018.5.PEDS1859
 55. Amorim JA, Remigio DS, Damazio Filho O, de Barros MA, Carvalho VN and Valença MM. **Intracranial subdural hematoma post-spinal anesthesia: report of two cases and review of 33 cases in the literature.** *Rev Bras Anesthesiol* 2010;60(6):620-629, 344-629 Doi:10.1016/S0034-7094(10)70077-5
 56. Perez Perez A, Calvo Porqueras B, Porta Etessam J and Jorquera Moya M. **Cerebrospinal fluid hypotension as a cause of cortical vein thrombosis.** *Neurologia* 2016;31(9):648-649 Doi:10.1016/j.nrl.2014.10.004
 57. Patel R, Urits I, Orhurhu V, Orhurhu MS, Peck J, Ohuabunwa E, . . . Viswanath O. **A Comprehensive Update on the Treatment and Management of Postdural Puncture Headache.** *Curr Pain Headache Rep* 2020;24(6):24 Doi:10.1007/s11916-020-00860-0
 58. Tonnelet R, Colnat-Coulbois S, Mione G, Richard S, Bouaziz H, Audibert G, . . . Braun M. **Successful Treatment of Spontaneous Intracranial Hypotension by Plugging the Cerebrospinal Fluid Leak with Percutaneous Cyanoacrylate Injection: A Report of 2 Cases.** *World Neurosurg* 2016;91:390-398 Doi:10.1016/j.wneu.2016.04.051



59. Janssen I, Gempt J, Gerhardt J, Meyer B and Ryang YM. **Treatment strategy for cerebral hypotension caused by spontaneous cerebrospinal fluid leaks.** *Acta Neurochir (Wien)* 2016;158(2):273-278 Doi:10.1007/s00701-015-2653-8
60. Vasdev GM and Chantigian RC. **Pneumocephalus following the treatment of a postdural puncture headache with an epidural saline infusion.** *J Clin Anesth* 1994;6(6):508-511 Doi:10.1016/0952-8180(94)90094-9
61. Stevens DS and Peeters-Asdourian C. **Treatment of postdural puncture headache with 'epidural dextran patch'.** *Reg Anesth* 1993;18(5):324-325, <http://www.ncbi.nlm.nih.gov/pubmed/7505607>