



Propranolol: A migraine prophylactic since the 1960s

Wallyson Pablo de Oliveira Souza¹, Yasmine Maria Leócido Fortes², Adriana de Almeida Soares²,
Raimundo Pereira Silva-Néto²

¹Federal University of the Parnaíba Delta, Parnaíba, Piauí, Brazil

²Graduate Program in Biomedical Sciences of the Federal University of the Parnaíba Delta, Parnaíba, Piauí, Brazil



Wallyson Pablo de Oliveira Souza
pablo_wallyson@hotmail.com

Edited by:

Marcelo Moraes Valença

Keywords:

Beta-blocker
Propranolol
Migraine
Prophylaxis

Abstract

Introduction

Propranolol was the first non-selective beta-adrenergic blocker to be developed. Initially it was used in the treatment of cardiovascular diseases, but since the 60's it has been used in the prevention of migraine.

Objective

The objective of this study was to know the history of propranolol and its use as a migraine prophylactic.

Methods

This study was an integrative literature review using articles with historical data on propranolol, from its origin in cardiology to its indication in the preventive treatment of migraine.

Results

Propranolol was described in 1962 for the treatment of cardiovascular diseases. In the same decade, it was prescribed for the preventive treatment of migraine and, recently, included in the consensus of the Brazilian Headache Society.

Conclusion

Although propranolol was initially synthesized for the treatment of heart disease, it has proved to be an effective drug in preventing migraine attacks.

Submitted: January 6, 2023.

Accepted: February 9, 2023.

Published Online: March 14, 2023



The origin of propranolol

Propranolol was the first non-selective beta-adrenergic blocker to be developed. It was synthesized from the insertion of an oxymethylene group in the structure of pronethalol, in 1962, by the Scottish pharmacologist James Whyte Black (1924-2010) and commercialized in 1964.¹⁻⁵

It is particularly indicated in the treatment of cardiovascular diseases such as systemic arterial hypertension, angina pectoris, hypertrophic obstructive cardiomyopathy, cardiac arrhythmias, hypertrophic subaortic stenosis and mitral valve prolapse and prevention of myocardial infarction.⁶⁻⁸ However it is also used in the treatment of essential tremor and in the prophylaxis of migraine.⁹

In the treatment of cardiovascular diseases, its mechanism of action is known. It blocks beta-adrenergic receptors, inhibiting sympathetic stimulation. This results in reduced resting heart rate, cardiac output, systolic and diastolic blood pressure, and reflex orthostatic hypotension.¹⁰

Propranolol should not be used in patients with cardiac decompensation, sinus dysfunction syndrome or second and third degree heart block, severe bradycardia, history of bronchospasm or bronchial asthma, chronic obstructive pulmonary disease, metabolic acidosis, diabetes and Prinzmetal's angina.

Who was Sir James Whyte Black

James Whyte Black was a Scottish pharmacologist who was born on 14 June 1924 and died on 23 March 2010, aged 85. He was professor of physiology at the University of Glasgow in Scotland; and Analytical Pharmacology at the Rayne Institute at King's College London Medical School, London.¹¹

Black was responsible for discovering two drugs: the first beta-blocker (propranolol), in 1962; and the first selective histamine H2 antagonist for the treatment of stomach ulcers (cimetidine), in 1975.¹¹

His research and discoveries developed his deep fascination with receptor theory as the foundation of pharmacology and drug discovery. He won the Nobel Prize in Medicine in 1988 for his method of drug invention, which consisted of building molecules around the structure of a natural chemical activator of a pathway involved in the etiology of a disease.³

The first prescriptions for migraine

Propranolol was first prescribed as a migraine prophylactic in the USA in 1968 by John Graham. Shortly afterwards, he stopped using it because that drug had not been approved by the Food and Drug Administration (FDA) to prevent migraine attacks.¹² A few years later, Graham performed a double-blind, placebo-controlled study and suggested that propranolol to be effective in preventing migraine.¹³

Still in 1968, Edgard Raffaelli Júnior, a disciple of John Graham, was the first physician in Latin America to prescribe propranolol as a preventive drug for migraine. He used propranolol on himself from 1968 until 1976, when he had the last headache attack, of short duration and mild intensity.^{12,14} Like Raffaelli, most headache pioneers tested drugs on themselves due to the lack of available options, as seen in the first publications in 1968.¹⁵

At that time, in Brazil, this drug had not been approved by the National Health Surveillance Agency for patients with migraine. As a result, Raffaelli faced problems with the prescription of propranolol because there was no reference to its indication for headache in the package insert. Some patients refused to use it because they did not have high blood pressure or heart disease.^{12,14}

In the early 1970s, studies began to appear demonstrating the effectiveness of propranolol in preventing migraines¹⁶⁻²⁰, but as it was not included in the package leaflet, Raffaelli decided to formulate all of his prescriptions. At that time, compounding pharmacies in São Paulo (Brazil) only prepared magistral formulas (those that appeared in medical books). Raffaelli was the first Brazilian doctor to create non-magisterial formulas.¹²

Propranolol and migraine

Prophylactic treatment of migraine came from Sicuteri's studies when he attributed to serotonin a prominent role in its genesis.²¹ Drugs that interfered with serotonin metabolism were tested. Initially, methysergide and pizotifen; then propranolol and amitriptyline.

Although propranolol has been used for migraine prophylaxis for over 50 years¹², only in 2002, Brazilian Headache Society appointed an ad hoc Committee with the purpose of establishing a consensus on the prophylactic treatment of migraine and of preparing recommendations to be disseminated among physicians. This Consensus recommended the use of tricyclic antidepressants, calcium channel blockers, serotonin antagonists, antiepileptic drugs



and beta-adrenergic blockers, including propranolol.²²

Propranolol has been widely used in migraine prophylaxis as a first-line medication.^{23,24} Despite the existence of consensus and guidelines that guide its prescription, many Brazilian doctors do not know how to prescribe this medication. They usually prescribe propranolol, wrongly, once a day or in inappropriate doses or even in long-term preparations that are not effective. The recommended dosage of propranolol ranges from 40 mg to 240 mg, two or three times a day, orally. It should start with low doses that can be gradually increased, according to the patient's response.²⁵⁻²⁷

Its exact mechanism of action in migraine prevention is not fully understood, but it lacks intrinsic sympathomimetic activity. Propranolol is believed to inhibit nitric oxide production by blocking kainite-induced currents and to have a synergistic effect on N-methyl-D-aspartate (NMDA) blockers, which reduce neuronal activity and have membrane-stabilizing properties.²⁸

Other authors have described migraine treatments in depth, including propranolol. The proposed mechanisms that could be cited are the inhibition of noradrenaline release due to the blockade of prejunctional beta-receptors; the reduction of firing rates at the locus ceruleus, the most important adrenergic nucleus in the brain; and the reduction of the noradrenaline synthesis through the action at the enzyme tyrosine-hydroxylase. In addition, an action downregulating 5HT_{2b} and c receptors is also a suggested effect.²⁹

Conclusion

Although propranolol was initially synthesized for the treatment of heart disease, it has proved to be an effective drug in preventing migraine attacks.

Wallyson Pablo de Oliveira Souza
<https://orcid.org/0000-0003-3122-9484>
 Yasmine Maria Leóido Fortes
<https://orcid.org/0000-0001-9642-0330>
 Adriana de Almeida Soares
<https://orcid.org/0000-0001-7002-6458>
 Raimundo Pereira Silva-Néto
<https://orcid.org/0000-0002-2343-9679>

Contribution authors: All authors had the same contribution.

Funding: No

Conflict of interests: The authors report no conflict of interest.

References

- Black JW and Stephenson JS. **Pharmacology of a new adrenergic beta-receptor-blocking compound (Nethalide).** *Lancet* 1962;2(7251):311-314 Doi:10.1016/s0140-6736(62)90103-4
- Black O, Gerskowitch SV, Hull RA and Shankley NP. **The pharmacological toolmaker's rational approach to drug design: An appreciation of Sir James Black.** *Trends Pharmacol Sci* 1988;9:435-437 Doi:10.1016/0165-6147(88)90132-0
- Stapleton MP. **Sir James Black and propranolol. The role of the basic sciences in the history of cardiovascular pharmacology.** *Tex Heart Inst J* 1997;24(4):336-342.
- Srinivasan AV. **Propranolol: A 50-year historical perspective.** *Ann Indian Acad Neurol* 2019;22(1):21-26 Doi:10.4103/aian.AIAN_201_18
- Dornhorst AC and Robinson BF. **Clinical pharmacology of a beta-adrenergic-blocking agent (Nethalide).** *Lancet* 1962;2(7251):314-316 Doi:10.1016/s0140-6736(62)90104-6
- Prichard BN. **Propranolol in the treatment of angina: a review.** *Postgrad Med J* 1976;52(Suppl 4):35-41.
- Lambert DM. **Effect of propranolol on mortality in patients with angina.** *Postgrad Med J* 1976;52(Suppl 4):57-60.
- Rabkin R, Stables DP, Levin NW and Suzman MM. **The prophylactic value of propranolol in angina pectoris.** *Am J Cardiol* 1966;18:370-383.
- Silva-Néto RP, Barbosa JMS and Almeida K. **Analysis of the package inserts of migraine prophylactics.** *Headache Medicine* 2010;1(1):12-16.
- Fitzgerald JD. **Beta-blockade and mechanisms of disease.** *Postgrad Med J* 1976;52(Suppl 4):184-190.
- Ganellin R and Duncan W. **Obituary: James Black (1924–2010).** *Nature* 2010;464:1292 Doi:10.1038/4641292a
- Silva-Néto RP. **Quem foi Edgard Raffaelli Júnior.** *Migrâneas cefaléias* 2006;9(4):152-158.
- Malvea BP, Gwon N and Graham JR. **Propranolol prophylaxis of migraine.** *Headache* 1973; 12(4): 163-167. doi: 10.1111/j.1526-4610.1973.hed1204163.x
- Silva-Néto RP. **The study of headache in the 1950s in Latin America by Edgard Raffaelli Júnior (1930-2006).** *Headache* 2015;55(5):713-717 Doi:10.1111/head.12578
- Hall GH. **Treatment of migraine with propranolol.** *Lancet* 1968;2(7578):1139 Doi:10.1016/s0140-6736(68)91607-3
- Weber RB and Reinmuth OM. **The treatment of migraine with propranolol.** *Neurology* 1972;22(4):366-369



- Doi:10.1212/wnl.22.4.366
17. Malvea BP, Gwon N and Graham JR. **Propranolol prophylaxis of migraine.** *Headache* 1973;12(4):163-167 Doi:10.1111/j.1526-4610.1973.hed1204163.x
 18. Borgesen SE, Nielsen JL and Moller CE. **Prophylactic treatment of migraine with propranolol. A clinical trial.** *Acta Neurol Scand* 1974; 50(5): 651-656 Doi:10.1111/j.1600-0404.1974.tb02810.x
 19. Ludvigsson J. **Propranolol used in prophylaxis of migraine in children.** *Acta Neurol Scand* 1974;50(1):109-115 Doi:10.1111/j.1600-0404.1974.tb01350.x
 20. Wideroe TE and Vigander T. **Propranolol in the treatment of migraine.** *Br Med J* 1974;2(5921):699-701 Doi:10.1136/bmj.2.5921.699
 21. Sicuteri F. **Prophylactic and therapeutic properties of 1-methylsergic acid butanolamide in migraine.** *Int Arch Allergy* 1959;15:300-307 Doi:10.1159/000229055
 22. Brazilian Headache Society. **Recommendations for prophylactic treatment of migraine: Consensus of the Sociedade Brasileira de Cefaléia.** *Arq Neuropsiquiatr* 2002; 60(1): 159-169 Doi: 10.1590/s0004-282x2002000100030
 23. Danesh A, Gottschalk PCH. **Beta-blockers for migraine prevention: A review article.** *Curr Treat Options Neurol* 2019;21(4):20 Doi:10.1007/s11940-019-0556-3
 24. Eigenbrodt AK, Ashina H, Khan S, Diener HC, Mitsikostas DD, Sinclair AJ, ... and Ashina M. **Diagnosis and management of migraine in ten steps.** *Nat Rev Neurol* 2021;17(8):501-514 Doi:10.1038/s41582-021-00509-5
 25. Loder E, Burch R and Rizzoli P. **The 2012 AHS/AAN guidelines for prevention of episodic migraine: a summary and comparison with other recent clinical practice guidelines.** *Headache* 2012;52(6):930-945 Doi:10.1111/j.1526-4610.2012.02185
 26. Cortelli P. **Low doses of propranolol are effective as migraine prophylaxis.** *Cephalalgia* 1990;10(3):153 Doi:10.1046/j.1468-2982.1990.1003153.x
 27. Pascual J, Polo JM and Berciano J. **The dose of propranolol for migraine prophylaxis. Efficacy of low doses.** *Cephalalgia* 1989;9(4):287-291 Doi:10.1046/j.1468-2982.1989.0904287.x
 28. Ramadan NM. **Prophylactic migraine therapy: mechanisms and evidence.** *Curr Pain Headache Rep* 2004;8(2):91-95 Doi:10.1007/s11916-004-0022-z
 29. Krymchantowski AV. **Conduas em Cefaleia: Avaliação e Tratamento.** São Paulo: Lippincott Williams & Wilkins, 2008:45-46.