



Temporomandibular disorder changes cervical muscle strength in young adults?

Geovanna Cristina Pereira Alves¹, Josmayara da Silva Caldas¹; Thayllane Costa Cardoso¹, Camila Vitória de Moraes Costa¹, Rodrigo Sousa Andrade¹, Matheus Rafael Feques Ferreira Nogueira¹, Ryan Pinheiro Castro¹, João Pedro da Fonseca de Paula¹, Gustavo Weyber Pereira Alves², Maria Claudia Gonçalves³

¹Student of the Physiotherapy course at CEUMA, University, Sao Luís, Maranhão, Brazil

²Doctor graduated from the Federal University of Maranhão, Sao Luís, Maranhão, Brazil

³Professor of the postgraduate Program in Environment and the Undergraduate course in Physiotherapy at CEUMA University, Sao Luís, Maranhão, Brazil

Introduction

Literature indicates relationships between temporomandibular disorder (TMD) pain and disability related to the cervical spine, but studies evaluating the strength of cervical muscles, especially in young adults, are still scarce.

Objectives

To evaluate cervical muscle strength in young adults with and without temporomandibular disorder.

Materials and Methods

This study included young adults aged 18 to 25 years from a private university, both with and without a diagnosis of TMD, and in generally good health. Individuals with neck/head trauma; disc herniation and degeneration; systemic diseases such as fibromyalgia, lupus; pregnancy; or anesthetic block in the last 3 months were excluded. Those scoring >50 points on the Fonseca Anamnestic Index questionnaire were considered to have TMD. The Neck Disability Index was used to assess the level of disability related to cervical spine pain. Cervical muscle strength of the flexors and extensors was measured by maximum isometric voluntary contraction using a handheld dynamometer, model 2201163, Lafayette, IN, USA. A significance level of <0.05 was adopted.

Results

A total of n=64 individuals were evaluated, with n=30 in the TMD group, averaging 20 to 22 years of age, with moderate TMD being the most observed (n=34, 53%). A reduction in muscle strength was identified in the TMD group compared to the non-TMD group for all movements, with a significant difference for the flexion movement (p<0.05).

Conclusion

Individuals with TMD show lower muscle strength for all cervical spine movements compared to those without the condition, with this difference being significant for the flexion movement.