



Case Report

"I feel an irresistible urge to bang my head against the wall": a case series highlighting agitation and severe pain in cluster headache

Marcelo Moraes Valença^{1,2}, Rita Santana dos Reis¹, Juliana Ramos de Andrade¹

¹Federal University of Pernambuco, Recife, Pernambuco, Brazil

²Hospital Esperança, Recife, Pernambuco, Brazil



Marcelo Moraes Valença
mmvalenca@yahoo.com.br

Edited by:

Mário Fernando Pietro Peres

Keywords:

Cluster headache
Psychomotor agitation
Head-banging
Self-injurious behavior
trigeminal autonomic cephalalgia

Introduction

Paroxysmal cranial sensations without pain are infrequently described and remain poorly classified in current headache nosology. While disorders such as primary stabbing headache and epicrania fugax produce brief, sharp pain, non-painful paroxysms of the scalp are rare and understudied. We report a case of recurrent, brief, non-painful thumping (or "blow") sensations at the vertex, persisting for seven years, to broaden awareness of atypical cranial sensory phenomena.

Case Report

A 60-year-old woman presented with a 7-year history of recurrent, spontaneous sensations she described as a "thump" or "blow" localized to the vertex. Individual episodes lasted ~2 seconds and recurred approximately once monthly. The patient denied any associated pain, auditory phenomena, visual disturbance, dizziness, nausea, or aura. Episodes occurred in diverse positions (lying, sitting, walking) and were not precipitated by head movement or Valsalva maneuvers. During events she briefly stopped activity due to a transient fear of imbalance but did not fall. Medical history was unremarkable for hypertension, diabetes, tobacco use, trauma, or recent illness. Neurological examination was normal. No focal deficits, post-ictal signs, or autonomic features were observed.

Comment and Conclusion

This presentation does not fit established ICHD-3 entities and likely represents a benign, focal paroxysmal sensory phenomenon localized to the scalp. Differential diagnoses include atypical primary sensory paroxysm, focal sensory epileptic discharge, and brief peripheral neuropathic events; however, the stereotyped, nonpainful, ultra-brief nature and long stability favor a benign primary sensory paroxysm. Reporting such cases can help refine classification and stimulate research into the pathophysiology of non-painful cranial paroxysms.

Received: September 7, 2025

Revised: September 20, 2025

Accepted: September 28, 2025



Introduction

Cluster headache is a primary headache disorder characterized by excruciating, unilateral pain, often accompanied by autonomic symptoms such as conjunctival injection, lacrimation, nasal congestion, rhinorrhea, eyelid oedema, forehead and facial sweating, miosis and ptosis (1-3).

One striking behavioral phenomenon frequently observed in these patients is intense psychomotor agitation (4), including an irresistible urge to bang one's head against a hard surface, such as a wall (4, 5).

This case series presents three patients whose extreme reactions to pain included the wish to engage in head-banging behavior, highlighting the profound suffering associated with cluster headache.

Case Report

Case 1

A 34-year-old man presented with a two-year history of severe, unilateral headaches localized to the right orbital and temporal regions. The attacks occurred in clusters lasting 3–4 months, followed by periods of remission. Based on clinical features and periodicity, the diagnosis of episodic cluster headache was established. The attacks occurred twice daily, typically lasting 60 to 90 minutes. During the crisis, the patient would pace the room, scream, and repeatedly want to hit his head against the wall in desperation. He described the pain as a "red-hot iron" being driven through his eye.

His neurological examination was normal. Brain MRI and MR angiography excluded structural or vascular causes. The diagnosis of episodic cluster headache was made according to ICHD-3 criteria. He was started on verapamil (240 mg/day) and high-flow oxygen during attacks. After up-titration to 480 mg/day, the frequency of attacks decreased significantly.

Case 2

A 38-year-old woman reported left-sided periorbital pain attacks with autonomic signs, including lacrimation and rhinorrhea, lasting approximately 45 minutes. The attacks occurred once to three times daily during active periods, with pain so intense she felt compelled to strike her head against a desk or wall. She reported a sense of helplessness and described the behavior not as suicidal, but as an impulsive response to unbearable pain.

The patient had no prior psychiatric history. Imaging studies were unremarkable. She was diagnosed with cluster headache and treated with a transitional corticosteroid course and prophylactic topiramate (100

mg/day). Oxygen therapy provided partial relief during attacks. After three weeks of treatment, the episodes became less frequent, and the need for self-injurious behavior dissipated.

Case 3

A 40-year-old man presented with a 10-year history of chronic cluster headache. His attacks occurred multiple times daily without significant remission periods. He described the pain as "inhuman" and acknowledged having banged his head against the floor or wall on several occasions during attacks. He explained that this behavior gave him a sense of control and was a way to express the unbearable inner torment.

He had previously failed several prophylactic treatments. At the time of evaluation, he was using sumatriptan injections with partial relief. He was started on high-dose verapamil (up to 720 mg/day) with close ECG monitoring. Occipital nerve block was performed as an adjunctive therapy. The patient reported a reduction in attack frequency and intensity, and the behavioral agitation was markedly attenuated.

Discussion

The urge to bang one's head against the wall during a cluster headache attack is a commonly reported behavior and reflects the excruciating intensity of the pain as well as the associated psychomotor agitation experienced during the crisis. This phenomenon can be explained by several neurobiological and psychological mechanisms:

Extreme pain and self-stimulatory behavior

Cluster headache is often described as one of the most severe types of pain known in clinical medicine. In the face of such overwhelming pain, the brain may attempt to redirect or modulate the sensation through external physical stimuli (5, 6). Hitting the head against a hard surface or applying pressure may be an unconscious attempt to: "Replace" the internal pain with an external stimulus that feels more controllable; Activate descending pain inhibitory pathways, such as the release of endorphins or alternative sensory input via spinothalamic circuits (7).

Motor agitation as a hallmark of the disorder

Unlike migraine, where patients typically seek silence, darkness, and rest; cluster headache sufferers often exhibit severe agitation, restlessness, or even self-injurious behavior during attacks (8). This pattern is thought to reflect the desperation and inability to find relief, rather than any intentional desire to harm oneself.



Hypothalamic and limbic system activation

Functional neuroimaging studies have shown activation of the posterior hypothalamus during cluster attacks (9-11). This region plays a key role not only in autonomic regulation but also in primitive emotional and behavioral responses, including aggression, impulsivity, and motor arousal (12-18). Thus, the combination of unbearable pain with hypothalamic-limbic activation may lead to disorganized motor behaviors such as pacing, shouting, or head banging.

Desperate attempt to find relief

Many patients report that no position or environment relieves the pain and describe a sense of utter hopelessness. Striking the head may be a desperate attempt to interrupt or counteract the internal torment, even if momentarily or symbolically (19).

Conclusion

This case series illustrates a dramatic but under-recognized behavioral manifestation of cluster headache: the urge to bang the head during attacks. It reflects both the severity of the pain and the neurobiological activation of structures involved in emotion and motor behavior. Clinicians should be aware of this symptom, as it underscores the importance of early diagnosis, comprehensive treatment, and compassionate care to reduce patient suffering and prevent such extreme responses.

Agitation alone is so strongly associated with cluster headache that its presence is sufficient to fulfill the ICHD-3 diagnostic criteria for cluster headache, even in the absence of any autonomic features (20).

References

- de Freitas Dias B, Robinson CL, Villar-Martinez MD, Ashina S, Goadsby PJ. Current and Novel Therapies for Cluster Headache: A Narrative Review. *Pain Ther.* 2025;14(2025):1-19. Doi: 10.1007/s40122-024-00674-7
- Xu XH, Li YM, Ren LN, Xu XF, Dai YL, Jin CQ, Yang RR. Cluster headache: understandings of current knowledge and directions for whole process management. *Front Neurol.* 2024;15(2024):1456517. doi: 10.3389/fneur.2024.1456517.
- Burish M. Cluster Headache, SUNCT, and SUNA. *Continuum (Minneapolis, Minn).* 2024;30(2024):391-410. Doi: 10.1212/CON.0000000000001411.
- Torelli P, Manzoni GC. Behavior during cluster headache. *Curr Pain Headache Rep.* 2005;9(2005):113-9. Doi: 10.1007/s11916-005-0048-x.
- Luerding R, Henkel K, Gaul C, Dresler T, Lindwurm A, Paelecke-Habermann Y, et al. Aggressiveness in different presentations of cluster headache: Results from a controlled multicentric study. *Cephalalgia.* 2012;32(2012):528-36. Doi: 10.1177/0333102412443336.
- Yulug B, Hanoglu L, Schabitz WR. Headache and spontaneous glabellar ecchymosis: More than a self-injury behavior? *Psychiatry and Clinical Neurosciences.* 2014;69(2014):124-. Doi: 10.1111/pcn.12220.
- Torelli P, Manzoni GC. Behavior during cluster headache. *Current Pain and Headache Reports.* 2005;9(2005):113-9. Doi: 10.1007/s11916-005-0048-x.
- Bertotti G, Fernandez-Ruiz V, Roldan-Ruiz A, Lopez-Moreno M. Cluster Headache and Migraine Shared and Unique Insights: Neurophysiological Implications, Neuroimaging, and Biomarkers: A Comprehensive Review. *J Clin Med.* 2025;14(2025). Doi: 10.3390/jcm14072160.
- Abagnale C, Di Renzo A, Giuliani G, Sebastianelli G, Casillo F, Ziccardi L, et al. MRI-based analysis of the microstructure of the thalamus and hypothalamus and functional connectivity between cortical networks in episodic cluster headache. *J Headache Pain.* 2025;26(2025):12. Doi: 10.1186/s10194-024-01920-1.
- Chiffi D, Di Renzo A, Giuliani G, Abagnale C, Altieri M, Sebastianelli G, et al. Assessment of hypothalamic macrostructure in episodic cluster headache: a volumetric segmentation MRI study. *Radiol Med.* 2025;130(2025):1483-8. Doi: 10.1007/s11547-025-02041-8.
- Coppola G, Abagnale C, Sebastianelli G, Goadsby PJ. Pathophysiology of cluster headache: From the trigeminovascular system to the cerebral networks. *Cephalalgia.* 2024;44(2024):3331024231209317. Doi: 10.1177/03331024231209317.
- Bai F, Huang L, Deng J, Long Z, Hao X, Chen P, et al. Prelimbic area to lateral hypothalamus circuit drives social aggression. *iScience.* 2023;26(2023):107718. Doi: 10.1016/j.isci.2023.107718.
- Rizzi M, Gambini O, Marras CE. Posterior hypothalamus as a target in the treatment of aggression: From lesioning to deep brain stimulation. *Handb Clin Neurol.* 2021;182(2021):95-106. Doi: 10.1016/B978-0-12-819973-2.00007-1.
- Vilela-Filho O. Commentary: Amygdala and Hypothalamus: Historical Overview With a Focus on Aggression. *Neurosurgery.* 2019;85(2019):E1-E3. Doi: 10.1093/neuros/nyz045.
- Gouveia FV, Hamani C, Fonoff ET, Brentani H, Alho EJJ, de Moraes R, et al. Amygdala and Hypothalamus: Historical Overview With Focus on Aggression. *Neurosurgery.* 2019;85(2019):11-30. Doi: 10.1093/neuros/nyy635.
- Chang CH, Gean PW. The Ventral Hippocampus



- Controls Stress-Provoked Impulsive Aggression through the Ventromedial Hypothalamus in Post-Weaning Social Isolation Mice. *Cell Rep.* 2019;28(2019):1195-205 e3. Doi: 10.1016/j.celrep.2019.07.005.
17. Haller J. The Role of the Lateral Hypothalamus in Violent Intraspecific Aggression-The Glucocorticoid Deficit Hypothesis. *Front Syst Neurosci.* 2018;12(2018):26. Doi: 10.3389/fnsys.2018.00026.
 18. Hashikawa Y, Hashikawa K, Falkner AL, Lin D. Ventromedial Hypothalamus and the Generation of Aggression. *Front Syst Neurosci.* 2017;11(2017):94. Doi: 10.3389/fnsys.2017.00094.
 19. Vaughan R. My Own Private Purgatory: How Cluster Headaches Affect My Life. *Headache: The Journal of Head and Face Pain.* 2008;48(2008):1541-3. Doi: 10.1111/j.1526-4610.2008.01303.x.
 20. Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. *Cephalalgia.* 2018;38(2018):1-211. Doi: 10.1177/0333102417738202.

Marcelo Moraes Valença
<https://orcid.org/0000-0003-0678-3782>
 Rita Santana dos Reis
<https://orcid.org/0000-0002-9868-2411>
 Juliana Ramos de Andrade
<https://orcid.org/0000-0002-5445-8872>

Authors contribution: MMV, conceptualization, supervision, writing – review & editing; RSR, writing – review & editing; JRA, writing – original draft, writing – review & editing.

Conflict of interest: We declare no conflicts of interest.

Funding: We have not received significant financial support.