

The characteristic warning signs of headache associated with cerebral aneurysm

*I*n this issue a special attention is drawn to the possibility that a headache may be caused by an unruptured intracranial aneurysm, a vascular lesion with a high risk of causing a catastrophic intracranial hemorrhage⁽¹⁾. Up to 10% of the population will develop an intracranial aneurysm during their lifetime. As a result, its clinical manifestations, such as headache, need to be studied in depth⁽²⁾.

The literature is still scanty regarding the unruptured aneurysm, even though a considerable amount of information can be found on the headache induced by a rupture of an intracranial aneurysm⁽³⁾. An intracranial aneurysm may trigger a headache manifestation through different mechanisms, e.g. direct compression of pain-sensitive intracranial structures and sudden saccular expansion^(2,4).

Aneurysms sometimes trigger pain simulating primary headaches, such as tension-type headache, stabbing headache or cluster headache^(5,6). Thus, in a patient with a recent onset of headache, a headache beginning after the age of 50, or triggered by head movements or Valsalva's maneuver, or when the pain is located on the same side of the head (lock-in headache), an investigation must be conducted in order to rule out the possibility of a cerebral aneurysm being the cause⁽²⁾.

The readers will also find several other articles dealing with migraine and the triggering food factors, cross-cultural adaptation of the Brazilian Portuguese version of Quality of Life Headache-Youth (QLH-Y) Questionnaire and the relationship between fibromyalgia and Headache.

References

1. Valença MM. "Sit back, observe, and wait." Or is there a pharmacologic preventive treatment for cerebral aneurysms? *Neurosurg Rev.* 2013;36(1):1-9.
2. Valença MM, Andrade-Valença LPA, Martins C. Functional anatomy of headaches: circle of Willis aneurysms, third cranial nerve and pain. *Headache Medicine.* 2011;2(3):82-8.
3. Valença MM, Valença LPAA. Hemorragia subaracnóidea: quadro clínico, diagnóstico e tratamento. *Neurobiologia.* 2000;63:97-104.
4. Asano AGC, Silva WF, Valença MM. Cefaléia sentinela: sinal de alerta da hemorragia subaracnóidea por ruptura de aneurisma intracraniano. *Migrâneas & Cefaléias.* 2008;11:78-83.
5. Valença MM, Andrade-Valença LPA, Oliveira DA, Silva JC, Martins C. 2012. Headache Associated with Intracranial Aneurysms. In *Headaches: Causes, Treatment and Prevention*, edited by Pietro G. Gallo; Giovanna M. Giordano. New York: Nova Publishers. Pg 65-94.
6. Valença MM, Andrade-Valença LP, Martins C, de Fátima Vasco Aragão M, Batista LL, Peres MF, da Silva WF. Cluster headache and intracranial aneurysm. *J Headache Pain.* 2007;8(5):277-82.

Marcelo Moraes Valença
Editor-in-chief
Headache Medicine