



Neuroimaging evaluation in headache patients who have suffered a stroke or traumatic brain injury

Maria de Fátima Viana Vasco Aragão^{1,2}, Luziany Carvalho Araújo³, Marcelo Moraes Valença¹

¹Universidade Federal de Pernambuco, Recife, Brazil

²Centro Diagnóstico Multimagem, Recife, Brazil

³Department of Radiology, Hospital da Restauração, Recife, Brazil



Maria de Fátima Viana Vasco Aragão
fatima.vascoaragao@gmail.com

Edited by:

Juliana Ramos de Andrade

Abstract

In a medical emergency, the most urgent patients at significant risk of death are those with a cerebrovascular accident and those with traumatic brain injury. Many are admitted with diminished conscience status (coma) and focal neurological deficits. In the evaluation of these patients, neuroimaging is indispensable in order to identify the type of lesion and the location of the brain where it is located.

In the case of stroke, we can subdivide it into hemorrhagic and ischemic. Among hemorrhagic hemorrhages, we can mention (1) spontaneous intracerebral hematomas and (2) hemorrhages due to rupture of an intracranial aneurysm, with subarachnoid hemorrhage leading.

Patients with head trauma are critical; even those who arrive at the hospital alert and oriented can decrease their level of consciousness in a few hours due to an intracranial hematoma, edema, or cerebral contusion.

Thus, the availability of performing neuroimaging evaluations, using computed tomography and magnetic resonance imaging, or even digital angiography, is vital for continuous supervision of this type of patient. The exams often require repetition several times due to the rate of evolution of vascular lesions and after head trauma.

A warning sign in these types of patients is headache. In the intracranial aneurysmal rupture, we classically have the thunderclap headache, an explosive, sudden pain mentioned as the worst pain the individual has suffered in his or her life. The pericranium and some intracranial structures are sensitive to nociceptive stimuli, such as the dura mater, large arteries, and venous sinuses. The brain is relatively insensitive to painful stimuli.

This narrative review aims to inform the importance of neuroimaging assessment of patients with stroke and traumatic brain injury in an emergency department. In conclusion, a neuroimaging evaluation is paramount in addition to a neurological and physical examination of the critically ill patient with cerebrovascular disease or who has suffered a traumatic brain injury.

Keywords:

Stroke
Headache
Cerebral aneurysm
MRI
Computed tomography
Emergency

Received: May 5, 2022
Accepted: June 22, 2022



Introduction

In a medical emergency, the most critical patients at significant risk of death are those who have suffered a cerebrovascular accident and those with traumatic brain injury.¹⁻³ Many are admitted with diminished conscience status (coma) and focal neurological deficits. In the evaluation of these patients, neuroimaging is essential in order to identify the type of lesion and the location of the brain where it is located.^{4,5}

In the case of stroke, we can subdivide it into hemorrhagic and ischemic. Among hemorrhagic hemorrhages, we find (1) spontaneous intracerebral hematomas⁶⁻⁸ and (2) hemorrhages due to rupture of an intracranial aneurysm, with subarachnoid hemorrhage leading^{9,10}.

Patients with head trauma are critical; even those who arrive at the hospital alert and oriented can decrease their level of consciousness in a few hours due to an intracranial hematoma, edema, or cerebral contusion.^{11,12}

Thus, the availability of performing neuroimaging evaluations, using computed tomography and magnetic resonance imaging, or even digital angiography, is vital for continuous surveillance of this type of patient. The exam

often needs to be repeated several times due to the quality of evolution of vascular lesions and after head trauma.

A warning sign in these patients is headache.^{13,14} In the intracranial aneurysmal rupture, we classically have the thunderclap headache, an explosive, sudden pain mentioned as the worst pain the individual has suffered in his or her life.^{15,16} The pericranium and some intracranial structures are sensitive to nociceptive stimuli, such as the dura mater, large arteries, and venous sinuses. The brain is relatively insensitive to painful stimuli.

This narrative review aims to inform the importance of neuroimaging assessment of patients with stroke and traumatic brain injury.

Traumatic Brain Injury

Head trauma can affect the soft parts (scalp, pericranium, and bone), causing a localized or diffuse headache for a few days. In more severe cases, trauma can damage the brain parenchyma, causing tissue damage (contusion) and even the formation of hematomas (Figure 1), which leads to more intense and lasting headaches.¹⁷⁻¹⁹




Figure 1. (A) Bilateral frontoparietal chronic subdural hematoma, and (B) right frontoparietal epidural hematoma. (A) Non-contrast computed tomography of the skull in the axial plane shows bilateral frontoparietal subdural hematomas (arrows), which are characterized by biconcave collections, with heterogeneous density because they present bleeding areas in several phases, suggesting that they are multiloculated (the more recent the bleeding areas, the greater the hyperdensity). It determines moderate compression on the surface of the cerebral hemispheres with effacement of the cortical sulci. (B) Magnetic resonance imaging of another patient, FLAIR axial sequence, showing a right frontoparietal epidural hematoma characterized by a hyperintense biconvex collection in this sequence, determining moderate compression over the right pre- and post-central gyri.



Vascular Headache

Intracranial Aneurysms

Occur in 1-2% of the population and account for about 80-85% of non-traumatic subarachnoid hemorrhages.²⁰ Intracranial aneurysms are sporadically acquired lesions; however, a rare familial form has been linked to conditions such as autosomal dominant polycystic kidney disease, Marfan syndrome, Ehlers-Danlos syndrome type IV, fibromuscular dysplasia, moyamoya disease, sickle cell

disease, and arteriovenous malformations of the brain.²⁰ An important risk factor for an aneurysm is family history.²⁰

Non-traumatic Subarachnoid Hemorrhage

Subarachnoid hemorrhage accounts for 5.5% to 15% of strokes, usually resulting from ruptured cranial aneurysms (Figure 2). Headache is sudden and severe, often accompanied by nausea, vomiting, and photophobia.^{19, 21} It lasts between 2 and 3 weeks and is associated with a high mortality rate.




Figure 2. Subarachnoid hemorrhage due to rupture of intracranial aneurysm of the right posterior communicating artery. On the axial computed tomography image of the skull (A), subarachnoid hemorrhage is observed, located mainly in the suprasellar, perichiasmatic and ambient cisterns on the right, identifying hyperdensity of the acute bleeding (arrows) that is distributed through the cistern of the skull base (subarachnoid space), described above. The temporal horns (arrowhead) are well visualized as they are slightly enlarged, characterizing a discrete communicating hydrocephalus determined by subarachnoid bleeding. On digital angiography (B), saccular dilatation is observed at the origin of the right posterior communicating artery (arrow).

Arterial Dissection

Cranio-cervical arterial dissection is a leading cause of stroke in young and middle-aged adults, but it can occur at any age.²²

The classic clinical history is headache or unilateral neck pain, accompanied by partial Horner syndrome and followed by cerebral ischemia hours or days later. It may be spontaneous or traumatic, and may result from mild trauma (cough, cervical chiropractic).²³

The dissection results from the rupture of the intima with extravasation of blood into the vessel wall, creating a

“false lumen”. The intramural hematoma can expand making the artery irregular and stenotic or even completely occluded. The formation of dissecting pseudoaneurysms is common.²⁴

On non-contrast computed tomography, a hematoma on the vessel wall can be evidenced, characterized by a spontaneously hyperattenuating crescent-shaped area. On computed tomography angiography, an increase in the thickness of the dissected artery can be observed with an eccentric and narrow lumen surrounded by the mural thrombus and the intimal flap. On MRI, the T1 sequence with fat saturation is the best to characterize the dissection, in which the mural thrombus classically presents



hypersignal. In angioresonance, we can see irregularities in the walls of the dissected vessel, reduction or absence of blood flow signal.²³

Reversible Cerebral Vasoconstriction Syndrome

Reversible cerebral vasoconstriction syndrome, also called

Call-Fleming syndrome and spasm-associated headache, is characterized by severe headache with or without neurological symptoms and vasospasms, determining a vascular appearance in a "necklace of beads". It is more frequent in women and can be precipitated by pregnancy, drug use, sexual intercourse, etc. Disappear spontaneously in 1-3 months. (Figure 3).^{25,26}




Figure 3. Vascular cause of headache, probable basilar artery dissection or vasospasm. Male patient reports onset of intense headache during sexual intercourse. Digital angiography (A) and magnetic resonance angiography with 3D TOF technique (B) showing moderate reduction in caliber and parietal irregularities in the middle and lower third of the basilar artery. There is dominance of the left vertebral artery and hypoplasia of the right (anatomical variation). Digital angiography (C) and magnetic resonance angiography with 3D TOF technique (D) control, two months later, showing that the basilar artery has normal caliber and contours, with complete resolution of the image and the patient's symptoms.



Conclusion

In conclusion, a neuroimaging evaluation is paramount in addition to a neurological and physical examination of the critically ill patient with cerebrovascular disease or who has suffered a traumatic brain injury.

No conflict of interest

Financing

The work was financed with own resources.

Authors' contribution

MFVVA and LCA, wrote the article and the selection of the figures; MMV, the final edit of the manuscript.

Acknowledgement

We wish to express our thanks for the co-operation extended in the elaboration of this chapter to the neurosurgeons Suzana Serra Reis and Maria de Fátima Leal Griz from Recife, Pernambuco.

Maria de Fátima Viana Vasco Aragão
<https://orcid.org/0000-0002-2341-1422>
 Luziany Carvalho Araújo
<https://orcid.org/0000-0001-5072-8487>
 Marcelo Moraes Valença
<https://orcid.org/0000-0003-0678-3782>

References

- Gwarzo IH, Perez-Patron M, Xu X, Radcliff T and Horney J. **Traumatic Brain Injury Related Hospitalizations: Factors Associated with In-hospital Mortality among Elderly Patients Hospitalized with a TBI.** *Brain Inj* 2021;35(5):554-562 Doi:10.1080/02699052.2021.1890822
- Ehelepola ND, Ranasinghe TI, Prashanthi B and Bandara HM. **An unusual presentation of a stroke in a developing country: a case report.** *BMC Res Notes* 2017;10(1):69 Doi:10.1186/s13104-017-2378-2
- Griesbach GS, Kreber LA, Harrington D and Ashley MJ. **Post-acute traumatic brain injury rehabilitation: effects on outcome measures and life care costs.** *J Neurotrauma* 2015;32(10):704-711 Doi:10.1089/neu.2014.3754
- Wiesmann M and Bruckmann H. **[Diagnostic imaging of acute head and brain injuries].** *Radiologe* 1998;38(8):645-658 Doi:10.1007/s001170050405
- Brown AM, Twomey DM and Wong Shee A. **Evaluating mild traumatic brain injury management at a regional emergency department.** *Inj Prev* 2018;24(5):390-394 Doi:10.1136/injuryprev-2018-042865
- Rizos T, Dorner N, Jenetzky E, Sykora M, Mundiyanapurath S, Horstmann S, . . . Steiner T. **Spot signs in intracerebral hemorrhage: useful for identifying patients at risk for hematoma enlargement?** *Cerebrovasc Dis* 2013;35(6):582-589 Doi:10.1159/000348851
- Baye M, Hintze A, Gordon-Murer C, Mariscal T, Belay GJ, Gebremariam AA and Hughes CML. **Stroke Characteristics and Outcomes of Adult Patients in Northwest Ethiopia.** *Front Neurol* 2020;11(428) Doi:10.3389/fneur.2020.00428
- Xiao H, Yang Y, Xi JH and Chen ZQ. **Structural and functional connectivity in traumatic brain injury.** *Neural Regen Res* 2015;10(12):2062-2071 Doi:10.4103/1673-5374.172328
- Silver AJ, Pederson ME, Jr., Ganti SR, Hilal SK and Michelson WJ. **CT of subarachnoid hemorrhage due to ruptured aneurysm.** *AJNR Am J Neuroradiol* 1981;2(1):13-22, <https://www.ncbi.nlm.nih.gov/pubmed/6784546>
- Zhang XH, Zhao XY, Liu LL, Wen L and Wang GX. **Identification of ruptured intracranial aneurysms using the aneurysm-specific prediction score in patients with multiple aneurysms with subarachnoid hemorrhages- a Chinese population based external validation study.** *BMC Neurol* 2022;22(1):201 Doi:10.1186/s12883-022-02727-w
- Goodfellow MJ, Medina JA, Proctor JL, Xu S, Gullapalli RP, Rangghran P, . . . Fiskum G. **Combined Traumatic Brain Injury and Hemorrhagic Shock in Ferrets Leads to Structural, Neurochemical, and Functional Impairments.** *J Neurotrauma* 2022;Doi:10.1089/neu.2022.0102
- Ahn SH, Savarraj JP, Pervez M, Jones W, Park J, Jeon SB, . . . Choi HA. **The Subarachnoid Hemorrhage Early Brain Edema Score Predicts Delayed Cerebral Ischemia and Clinical Outcomes.** *Neurosurgery* 2018;83(1):137-145 Doi:10.1093/neuros/nyx364
- Quaid PT and Singman EL. **Post-traumatic headaches and vision: A review.** *NeuroRehabilitation* 2022;50(3):297-308 Doi:10.3233/NRE-228013
- Russo A, D'Onofrio F, Conte F, Petretta V, Tedeschi G and Tessitore A. **Post-traumatic headaches: a clinical overview.** *Neurol Sci* 2014;35 Suppl 1(153-156) Doi:10.1007/s10072-014-1758-9
- Tawk RG, Hasan TF, D'Souza CE, Peel JB and Freeman WD. **Diagnosis and Treatment of Unruptured Intracranial Aneurysms and Aneurysmal Subarachnoid Hemorrhage.** *Mayo Clin Proc* 2021;96(7):1970-2000 Doi:10.1016/j.



- mayocp.2021.01.005
16. Gilard V, Grangeon L, Guegan-Massardier E, Sallansonnet-Froment M, Maltete D, Derrey S and Proust F. **Headache changes prior to aneurysmal rupture: A symptom of unruptured aneurysm?** *Neurochirurgie* 2016;62(5):241-244 Doi:10.1016/j.neuchi.2016.03.004
 17. Parizel PM, Voormolen M, Van Goethem JW and van den Hauwe L. **Headache: when is neuroimaging needed?** *JBR-BTR* 2007;90(4):268-271, <https://www.ncbi.nlm.nih.gov/pubmed/17966243>
 18. Holle D and Obermann M. **The role of neuroimaging in the diagnosis of headache disorders.** *Ther Adv Neurol Disord* 2013;6(6):369-374 Doi:10.1177/1756285613489765
 19. Parizel PM, Makkat S, Van Miert E, Van Goethem JW, van den Hauwe L and De Schepper AM. **Intracranial hemorrhage: principles of CT and MRI interpretation.** *Eur Radiol* 2001;11(9):1770-1783 Doi:10.1007/s003300000800
 20. Ajiboye N, Chalouhi N, Starke RM, Zanaty M and Bell R. **Unruptured Cerebral Aneurysms: Evaluation and Management.** *ScientificWorldJournal* 2015;2015(954954) Doi:10.1155/2015/954954
 21. Kwon OK. **Headache and Aneurysm.** *Neuroimaging Clin N Am* 2019;29(2):255-260 Doi:10.1016/j.nic.2019.01.004
 22. Zhang FL, Guo ZN, Liu Y, Luo Y and Yang Y. **Dissection extending from extra- to intracranial arteries: A case report of progressive ischemic stroke.** *Medicine (Baltimore)* 2017;96(21):e6980 Doi:10.1097/MD.0000000000006980
 23. Rodallec MH, Marteau V, Gerber S, Desmottes L and Zins M. **Craniocervical Arterial Dissection: Spectrum of Imaging Findings and Differential Diagnosis.** *RadioGraphics* 2008;28(6):1711-1728 Doi:10.1148/rg.286085512
 24. Schwartz NE, Vertinsky AT, Hirsch KG and Albers GW. **Clinical and radiographic natural history of cervical artery dissections.** *J Stroke Cerebrovasc Dis* 2009;18(6):416-423 Doi:10.1016/j.jstrokecerebrovasdis.2008.11.016
 25. Calabrese LH, Dodick DW, Schwedt TJ and Singhal AB. **Narrative Review: Reversible Cerebral Vasoconstriction Syndromes.** *Annals of Internal Medicine* 2007;146(1):Doi:10.7326/0003-4819-146-1-200701020-00007
 26. Valenca MM, Valenca LP, Bordini CA, da Silva WF, Leite JP, Antunes-Rodrigues J and Speciali JG. **Cerebral vasospasm and headache during sexual intercourse and masturbatory orgasms.** *Headache* 2004;44(3):244-248 Doi:10.1111/j.1526-4610.2004.04054.x