



## Short Communication

# Why does ingestion of watermelon trigger headache attacks in patients with migraine?

Raimundo Pereira Silva-Néto<sup>1</sup> , Adriana Almeida Soares<sup>1</sup> , Luciano Silva Lopes<sup>2</sup> 

<sup>1</sup>Federal University of the Parnaíba Delta, Parnaíba, Brazil

<sup>2</sup>Federal University of Piauí, Teresina, Brazil



Raimundo Pereira Silva-Néto  
netoesperantina@terra.com.br

### Edited by:

Marcelo Moraes Valença

Several triggers can trigger a migraine attack, including food. By the way, food only triggers headache in migraine sufferers. The foods that most trigger headache attacks are these: cheese, chocolate, citrus fruits and some sweet fruits, such as watermelon.<sup>1</sup>

Foods are involved in several mechanisms in generating pain. They can lead to vasodilation (vascular effect), cause an increase in cytokines (inflammatory effect), activate peripheral glutamate receptors (effect on neuroreceptors), activate sympathetic and parasympathetic function (effect on the autonomic nervous system), and induce neuronal depolarization (cortical effect).<sup>2</sup>

Watermelon is involved in one of these pain-generating mechanisms. It is the main source of citrulline in nature. Citrulline is an amino acid that was first found in watermelon in 1930 and its name comes from *citrulus*, Latin name for watermelon (*Citrulus lanatus*).<sup>3</sup> Citrulline is responsible for the synthesis of arginine, the precursor of nitric oxide. In turn, nitric oxide is one of those responsible for generating pain.<sup>4</sup>

A recent experimental study carried out with two groups of volunteers, one with migraine and the other without migraine, showed that the ingestion of watermelon triggered headache attacks only in patients with migraine (23.7% vs. 0.0%;  $p=0.002$ ). The time interval between ingestion of watermelon and onset of headache was on average  $124.3 \pm 20.5$  minutes. In this study, serum levels of nitrite (the breakdown product of nitric oxide) were measured before and after ingestion of watermelon. There was an increase in serum levels of nitrite, compared to the baseline level ( $p < 0.001$ ).<sup>5</sup>

Submitted: July 14, 2023  
Accepted: November 27, 2023  
Published online: December 28, 2023



**Contribution authors:** All authors had the same contribution.

**Funding:** No

**Conflict of interests:** The authors report no conflict of interest.

Raimundo Pereira Silva-Néto

<http://orcid.org/0000-0002-2343-9679>

Adriana de Almeida Soares

<http://orcid.org/0000-0001-7002-6458>

Luciano da Silva Lopes

<http://orcid.org/0000-0002-7702-7258>

## References

1. Silva-Néto RP, Soares AA, Vasconcelos CAC and Lopes LS. **Watermelon and others plant foods that trigger headache in migraine patients.** *Postgrad Med* 2021;133(7):760-644 Doi:10.1080/00325481.2021.1922211.
2. Martin VT and Brinder V. **Diet and headache: Part 1.** *Headache* 2016;56(6):1543-1552 Doi:10.1111/head.12953.
3. Rimando AM and Perkins-Veazie PM. **Determination of citrulline watermelon rind.** *J Chromatogr A* 2005;1078(1-2):196-200 Doi:10.1016/j.chroma.2005.05.009.
4. Bailey SJ, Blackwell JR, Williams E, Vanhatalo A, Wylie LJ, Winward PG, ... and Jones AM. **Two weeks of watermelon juice supplementation improves nitric oxide bioavailability but not endurance exercise performance in humans.** *Nitric Oxide* 2016;59:10-20 Doi:10.1016/j.niox.2016.06.008.
5. Silva-Néto RP, Bezerra GL, Araújo NRA, Silva SF, Pereira SKS, Lima LKF, ... and Lopes LS.. **Migraine attacks triggered by ingestion of watermelon.** *Eur Neurol* 2023 2023;86(4):250-255 Doi:10.1159/000531286.