

Food avoidance among patients with headache

Alimentos que pacientes com cefaleia evitam

Stella Boreggio Machado¹
 Nayara Cavalcanti Ares⁻¹
 Claudio Scorcine²
 Yara Dadalti Fragoso³

¹ Universidade Metropolitana de Santos, Nutrition - Santos - SP - Brazil.

¹ Universidade Metropolitana de Santos, Physical Education - Santos - SP - Brazil.

¹ Universidade Metropolitana de Santos, Post Graduate Studies - Santos - SP - Brazil.

² MS & Headache Research, Research - Santos - SP - Brazil.

ABSTRACT

Objective: To assess food and drink avoidance among patients with headache by means of an online survey. **Methods:** Individuals with frequent headaches were invited to answer a Google Form questionnaire [https://form.jotformz.com/200233754863656]. The survey included gender, age and characteristics of headache. Dietary habits were assessed as the number of times the individual consumed certain foods, on a daily, weekly, or monthly basis. The participants could state up to three foods that they avoided for fear of headache attacks.

Results: 120 complete forms were received. Alcoholic beverages were the most frequent trigger factor, reported by 26.7% of the patients. 95.5% of the participants did not consume alcohol regularly. Cheese, caffeine and fat were also recognized as potential triggers of headaches. There was no standard profile of dietary triggers and, therefore, everyone has to be personally approached in this subject. **Conclusion:** The online survey confirmed that individual characteristics of headache were dietary triggers in half the participants. Alcohol was the most frequently mentioned trigger, followed by cheese, fat and caffeine.

Keywords: diet; headache; carbohydrates.

RESUMO

Objetivo: Avaliar quais comidas e bebidas pacientes com cefaleia evitam, usando um questionário online. **Métodos:** Pessoas com cefaleias frequentes foram convidadas a responder um questionário do Google Form [https://form.jotformz.com/200233754863656]. Esta pesquisa incluía sexo, idade, e características da cefaleia. Hábitos alimentares foram avaliados pelo número de vezes que o indivíduo ingeria certos alimentos de forma diária, semanal ou mensal. Os participantes podiam preencher até três tipos de alimento que evitassem com receio de desencadear crises de cefaleia. **Resultados:** A pesquisa recebeu 120 respostas completas. Bebidas alcoólicas foram o gatilho mais frequente, sendo referidas por 26,7% dos pacientes. Alcool não era consumido regularmente por 95,5% dos participantes. Queijo, cafeína e comidas gordurosas foram também reconhecidas como potenciais gatilhos de cefaleia. Não houve um perfil padrão de gatilhos dietéticos, portanto cada paciente deve ser abordado de maneira individual neste tema. **Conclusão:** A pesquisa online confirmou as características individuais dos gatilhos dietéticos de cefaleia em metade dos pacientes. Alcool foi o gatilho mais frequentemente relatado, seguido de queijo, gorduras e cafeína.

Descritores: dieta; cefaleia; carboidratos.

The association of dietary factors and primary headaches is controversial. Certain foods can trigger headache in up to 64% of patients, but not all the attacks and not all the time². The literature on this subject is conflicting since no mechanism for supporting the existence of a food-headache association “food-headache” has yet been established with adequate evidence³. Among the proposed mechanisms for the onset of headache attacks through dietary

triggers are the “amine hypothesis”, “allergy vasodilation”, “dysregulation of neurotransmitters involved in appetite” and “inflammatory diet”³.

Perhaps one of the best examples of the conflicting evidence on triggering foods relates to chocolate. Although eating chocolate is widely believed to trigger migraine attacks, the risk of having a migraine after doing this is as likely as after eating placebo⁴. Another confounding factor may be the masticatory trigger

*Correspondence

Yara Dadalti Fragoso
 E-mail: yara@bsnet.com.br

Received: April 24, 2020.
 Accepted: April 27, 2020.

DOI: 10.5935/2178-7468.20200013

for headache attacks: this could be misinterpreted as the food itself (for example, chewing red meat)⁵. Anxiety and anticipatory behavior can also play a role among patients who believe a certain food will trigger an attack⁶.

The objective of the present study was to assess headache patients' food avoidance and consumption using an online survey.

METHODS

This study was approved by the Ethics Committee at Universidade Metropolitana de Santos, SP, Brazil, under CAAE 17241719.1.0000.5509. Individuals with headache at least once a month over the last three months were invited to answer a Google Forms survey [https://form.jotformz.com/200233754863656].

The survey included gender, age and characteristics of headache. Dietary habits were assessed as the number of occasions on which the person consumed certain foods and drinks, on a daily, weekly, or monthly basis. Patients could fillup to three foods they spontaneously avoided for fearing of triggering headache attacks.

The Kolmogorov-Smirnov test assessed the sample normality, Student's t test was used for parametric variables and Pearson's correlation and the chi-square test were used for comparisons.

RESULTS

A total of 120 individuals answered the survey (95 women). Their average age was 36 years and 65.8% of them had presented headaches for four or more years. Migraine or probable migraine was identified in 104 subjects. The remaining 16 patients presented the characteristics of tension-type headache. Table 1 presents the list of foods that the patients avoided because they believed that these foods could induce headache attacks. Alcohol, greasy food, cheese and caffeine were the most cited triggers of headaches. Figure 1 presents

Table 1. Number (and percentage from n=120) of participants who spontaneously referred specific dietary components that could trigger headaches.

Food/drink	Number of patients (n)	%
Triggers?		
Yes	63	52.5%
Alcohol	32	26.6%
Bread	1	0.8%
Caffeine	9	7.5%
Cheese	9	7.5%
Chocolate	7	5.8%
Cured meats	4	3.3%
Egg	1	0.8%
Fat	15	12.5%
Nuts	3	2.5%
Red meat	2	1.6%
Salt	3	2.5%
Soda	3	2.5%
Spicy food	4	3.3%
Sweets	9	7.5%




Figure 1. Percentage of individuals with headache (n=120) who avoided consuming each of these foods or drinks. Note that avoidance of dietary factors is higher than that spontaneously cited by patients.

the frequencies of food consumption. In summary, alcoholic beverages, fizzy drinks, fruits, processed fruit juices, fish, prawns, soya products and cured meats were often avoided by these patients. Bread, cheese, natural fruit juices, beans, eggs, red meats, chocolates and coffee were frequently consumed by these patients. Only 13 patients (10.8%) reported five or more dietary triggers for their headaches that they never consumed.

There were no differences in food preference and/or avoidance regarding gender, age, frequency or type of headache. Caffeine and cheese, which were spontaneously cited as headache triggers by 10% of the patients, were among the five items most consumed by them. Cheese was consumed by 73% of the patients while black coffee was consumed by 77% of them. Canned fish and prawns were consumed by less than 10% of the patients, although none of them regarded these items as potential triggers.

DISCUSSION

The association between headaches (particularly migraine) and dietary components is complex and often misunderstood. Physicians frequently tell patients to avoid a list of standard foods and drinks that are not triggers for all headache sufferers. In addition, it is important to acknowledge that, beyond diet, other lifestyle changes may have a role in the therapeutic success of these patients⁷. Rather than implementing a standard list of foods and drinks that are "forbidden", identification of dietary triggers for each patient is ideal. This can be done with the help of food diaries, which are an inexpensive way to understand which foods and drinks may trigger headache attacks in that individual⁸.

It was interesting to observe that half the patients considered that at least one dietary component was a headache trigger. While alcohol was believed to trigger headache attacks by over a quarter of our patients, other dietary factors were remarkably different among the patients. For example, some participants could not tolerate cheese, while other ate it regularly without problems. This reinforces the idea that, like the pharmacological approach to headaches, a tailor-made dietary recommendation for each patient is necessary. While patients may give us details of their food avoidance, the biological mechanism



through which dietary triggers precipitate headache attacks remains obscure⁹.

Our study had limitations. It used a small sample of individuals who answering an online survey. The diagnosing their headache was not ideal, and the sample comprised a mixture of cases of migraine, probable migraine, and tension-type headache. However, the aim of this study was not to study any specific primary headache. There were no evaluations for micronutrients or the percentage of proteins, carbohydrates, and fat in the patients' diet. All of these factors will be addressed in future studies in our group.

CONCLUSION

The online survey used in this study confirmed the individual characteristics of headache dietary triggers. Alcohol, the most frequently reported trigger, affected 26% of the participants. Overall, half the patients had at least one food or drink that was associated with headaches.

ACKNOWLEDGEMENT

SB Machado received a Scientific Initiation Grant from the Brazilian National Council for Scientific and Technological Development (CNPq) for which the authors are grateful.

REFERENCES

1. Rockett FC, de Oliveira VR, Castro K, et al. Dietary aspects of migraine trigger factors. *Nutr Rev.* 2012; 70: 337-56. doi: 10.1111/j.1753-4887.2012.00468.x.
2. Zaeem Z, Zhou L, Dilli E. Headaches: a review of the role of dietary factors. *Curr Neurol Neurosci Rep.* 2016; 16: 101.
3. Razeghi Jahromi S, Ghorbani Z, Martelletti P, et al. Association of diet and headache. *J Headache Pain.* 2019; 20: 106. doi: 10.1186/s10194-019-1057-1
4. Lippi G, Mattiuzzi C, Cervellin G. Chocolate and migraine: the history of an ambiguous association. *Acta Biomed.* 2014; 85: 216-21.
5. Costa YM, Porporatti AL, Stuginski-Barbosa J, et al. Headache attributed to masticatory myofascial pain: clinical features and management outcomes. *J Oral Facial Pain Headache.* 2015; 29: 323-30. doi: 10.11607/ofph.1394.
6. Iliopoulos P, Damigos D, Kerezoudi E, et al. Trigger factors in primary headaches subtypes: a cross-sectional study from a tertiary centre in Greece. *BMC Res Notes.* 2015; 8: 393. doi: 10.1186/s13104-015-1390-7.
7. Slavin M, Ailani J. A clinical approach to addressing diet with migraine patients. *Curr Neurol Neurosci Rep.* 2017; 17: 17. doi: 10.1007/s11910-017-0721-6.
8. Sun-Edelstein C, Mauskop A. Foods and supplements in the management of migraine headaches. *Clin J Pain.* 2009; 25: 446-52. doi: 10.1097/AJP.0b013e31819a6f65.
9. Finocchi C, Sivori G. Food as trigger and aggravating factor of migraine. *Neurol Sci.* 2012; 33: S77-80. doi: 10.1007/s10072-012-1046-5.